

BEHAVIORAL COUNSELING INTERVENTIONS TO PROMOTE A HEALTHFUL DIET AND PHYSICAL ACTIVITY FOR CARDIOVASCULAR DISEASE PREVENTION IN ADULTS 2015	
Population	General adult population without a known diagnosis of hypertension, diabetes, hyperlipidemia, or cardiovascular disease (CVD)
Recommendation	<p>Although the correlation among healthful diet, physical activity, and the incidence of CVD is strong, existing evidence indicates that the health benefit of initiating behavioral counseling in the primary care setting to promote a healthful diet and physical activity is small. Clinicians may choose to selectively provide this service to patients rather than incorporating it into the care of all adults in the general population.</p> <p>Considerations: Clinicians may consider other risk factors for CVD, patient readiness for change, social support and community resources that support behavioral change, and other health care and preventive service priorities. Potential Harms: Harms may include the lost opportunity to provide other services with a greater health effect.</p> <p>Grade: C</p>
Risk Assessment	If an individual's risk for CVD is uncertain, there are several calculators and models available to quantify a person's 10-year risk for cardiac events, such as the Framingham-based Adult Treatment Panel III calculator (available at http://hp2010.nhlbi.nih.net/atp3iii/calculator.asp). Generally, persons with a 10-year risk greater than 20% are considered to be high-risk, those with a 10-year risk less than 10% are considered to be low-risk, and those in the 10% to 20% range are considered to be intermediate-risk.
Interventions	Medium- or high-intensity behavioral interventions to promote a healthful diet and physical activity may be provided to individual patients in primary care settings or in other sectors of the health care system after referral from a primary care clinician. In addition, clinicians may offer healthful diet and physical activity interventions by referring the patient to community-based organizations. Strong linkages between the primary care setting and community-based resources may improve the delivery of these services.
Balance of Harms and Benefits	The USPSTF concludes with moderate certainty that medium- or high-intensity primary care behavioral counseling interventions to promote a healthful diet and physical activity have a small net benefit in adult patients without CVD, hypertension, hyperlipidemia, or diabetes.
Other Relevant USPSTF Recommendations	The USPSTF has made recommendations on screening for carotid artery stenosis, coronary heart disease, high blood pressure, lipid disorders, obesity, and peripheral arterial disease. These recommendations are available at www.uspreventiveservicestaskforce.org .